

PCT/GIB/2004/007502

PA 1239208

THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE

United States Patent and Trademark Office

October 22, 2004

REC'D 16 DEC 2004

WIPO

PCT

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE UNDER 35 USC 111.

APPLICATION NUMBER: 60/512,200

FILING DATE: October 20, 2003

PRIORITY DOCUMENT

SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

By Authority of the
COMMISSIONER OF PATENTS AND TRADEMARKS



N. Williams
N. WILLIAMS
Certifying Officer

BEST AVAILABLE COPY

Mail Stop Provisional Patent Application

PTO/SB/16 (6-95)
Approved for use through 04/11/98. OMB 0651-0037
Patent and Trademark Office, U.S. DEPARTMENT OF COMMERCE

PROVISIONAL APPLICATION COVER SHEET

This is a request for filing a PROVISIONAL APPLICATION under 37 CFR 1.53 (c).
Docket Number 620-281 Type a plus sign (+) inside this box → +

INVENTOR(S)/APPLICANT(S)			
LAST NAME	FIRST NAME	MIDDLE INITIAL	RESIDENCE (CITY AND EITHER STATE OR FOREIGN COUNTRY)
OXFORD	Alexander	W.	Royston, United Kingdom
DAVIS	Richard	J.	Hitchin, United Kingdom
COLEMAN	Robert	A.	Royston, United Kingdom
CLARK	Kenneth	L.	Cambridge, United Kingdom
CLARK	David	E.	Harlow, United Kingdom
HARRIS	Neil	Victor	Harlow, United Kingdom
FENTON	Garry		Harlow, United Kingdom
HYND	George		Harlow, United Kingdom
STUTTLE	Keith Alfred	J.	Harlow, United Kingdom
SUTTON	Jonathan	M.	Harlow, United Kingdom

03917 U.S. PTO
60/512200
102003

TITLE OF THE INVENTION (280 characters)

EP, RECEPTOR ANTAGONISTS

CORRESPONDENCE ADDRESS

Direct all correspondence to:



Customer Number:

23117

Place Customer
Number Bar -
Label Here →

Type Customer Number here

ENCLOSED APPLICATION PARTS (check all that apply)



Specification

Number of Pages

59



Applicant claims "small entity" status.



"Small entity" statement attached.



Drawing(s)

Number of Sheets



Other (specify)

METHOD OF PAYMENT (check one)



A check or money order is enclosed to cover the Provisional filing fees (\$160.00)/(\$80.00)



The Commissioner is hereby authorized to charge any deficiency, or credit any overpayment, in the fee(s) filed, or asserted to be filed, or which should have been filed herewith (or with any paper hereafter filed in this application by this firm) to our Account No. 14-1140. A duplicate copy of this sheet is attached.

PROVISIONAL
FILING FEE
AMOUNT (\$)

160.00

The invention was made by an agency of the United States Government or under a contract with an agency of the United States Government.



No.



Yes, the name of the U.S. Government agency and the Government contract number are:

Respectfully submitted,
SIGNATURE

B. J. Sadoff

DATE

October 20, 2003

REGISTRATION NO.
(if appropriate)

36,663

TYPED or PRINTED NAME

B. J. Sadoff

PROVISIONAL APPLICATION FILING ONLY

Burden Hour Statement: This form is estimated to take .2 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Mail Stop Comments - Patents, Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450, and to the Office of Information and Regulatory Affairs, Office of Management and Budget (Project 0651-0037), Washington, DC 20503. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Mail Stop Provisional Patent Application, Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450.

784833

EP₄ RECEPTOR ANTAGONISTS

This invention relates to EP₄ receptor antagonists, pharmaceutical compositions comprising such compounds, and
5 the use of such compounds and compositions to treat various diseases.

Background to the invention

Prostanoids comprise prostaglandins (PGs) and thromboxanes
10 (Tx_s) and their receptors fall into five different classes (DP, EP, FP, IP and TP) based on their sensitivity to the five naturally occurring prostanoids, PGD₂, PGE₂, PGF_{2α}, PGI₂
and Tx_{A₂}, respectively (Coleman, R.A., Prostanoid Receptors.
IUPHAR compendium of receptor characterisation and
15 classification, 2nd edition, 338-353, ISBN 0-9533510-3-3, 2000). EP receptors (for which the endogenous ligand is PGE₂) have been subdivided into four types termed EP₁, EP₂, EP₃ and EP₄. These four types of EP receptors have been
cloned and are distinct at both a molecular and
20 pharmacological level (Coleman, R.A., 2000)

EP₄ antagonists have been shown to be useful in the treatment of pain, and in particular, in the treatment of primary headache disorders, which include migraines, and
25 secondary headache disorders, such as drug-induced headaches (WO 00/18405 and WO 01/72302). Dilation of the cerebral vasculature and the subsequent stimulation of pain stimulating, perivascular trigeminal sensory afferent nerves is recognised to play an important role in the
30 pathophysiology of migraine. A sterile inflammatory response, associated with activation of cyclooxygenase and the generation of PGE₂, is also implicated in the pathophysiology of migraine. PGE₂ levels have been shown to be raised during migraine attacks and PGE₂ contributes to

the pain of migraine by directly dilating cerebral arteries and by stimulating the release of vasoactive/pro-inflammatory peptides from the trigeminal nerves. These effects of PGE₂ are mediated in whole or in part by EP₄ receptors. Thus, by binding to and preventing the stimulation of EP₄ receptors, EP₄ antagonists may be used to treat the pain of migraine.

EP₄ antagonists may also be useful in treating a number of other conditions and diseases. For example, they may be used in:

- the treatment of pain associated with rheumatoid arthritis, osteoarthritis, rheumatoid spondylitis, gouty arthritis and juvenile arthritis;
- the treatment of musculoskeletal pain, lower back and neck pain, sprains and strains, neuropathic pain, sympathetically mediated pain, myositis, pain associated with cancer and fibromyalgia, pain associated with influenza or other viral infections, such as the common cold, rheumatic fever; pain associated with bowel disorders such as non-ulcer dyspepsia, irritable bowel syndrome; non-cardiac chest pain, pain associated with myocardial ischaemia, post-operative pain, headache, toothache and dysmenorrhea. Neuropathic pain syndromes include diabetic neuropathy, sciatica, non-specific lower back pain, multiple sclerosis pain, fibromyalgia, HIV-related neuropathy, post-herpetic neuralgia, trigeminal neuralgia and pain resulting from physical trauma;
- the treatment of inflammatory diseases including rheumatoid and osteoarthritis, psoriasis, dermatitis, retinitis, conjunctivitis, asthma, bronchitis, chronic obstructive pulmonary disease, inflammatory bowel disease, colitis, nephritis, gingivitis and hepatitis;
- the treatment of cancers including familial adenomatous


- polyposis, endometrial carcinoma, colorectal and cervical cancer;
- the treatment of bone disorders involving altered bone formation or resorption such as osteoporosis;
- 5 women's health for the treatment of myometrial and endometrial disorders;
- the treatment of gastrointestinal disease including diarrhoea;
- the treatment of immunological disorders such as autoimmune
- 10 disease, immunological deficiency diseases, organ transplantation and increasing the latency of HIV infection;
- the treatment of diseases of abnormal platelet function.
(e.g. occlusive vascular diseases);
- the preparation of a drug with diuretic properties to treat
- 15 or prevent various oedema, hypertension, premenstrual tension, urinary calculus, oliguria, hyperphosphaturia, mesangial proliferative glomerulonephritis, chronic renal failure or the like;
- the treatment of impotence or erectile dysfunction, and
- 20 female sexual dysfunction;
- the treatment of hair growth disorders;
- the treatment of sleep disorders such as narcolepsy and insomnia;
- the treatment of cardiovascular diseases and shock states
- 25 associated with hypotension (e.g. septic shock);
- the treatment of neurodegenerative diseases and for preventing neuronal damage following stroke, cardiac arrest, cardiopulmonary bypass, traumatic brain injury or spinal cord injury;
- 30 the treatment of tinnitus;
- the treatment of dependence; and
- the treatment of complications of diabetes.

Although EP₄ antagonists are known, it is desired to find

novel EP₄ antagonists, and in particular, EP₄ antagonists which are selective against other EP receptors, i.e. EP₁, EP₂ and EP₃.

5 Summary of the invention

A first aspect of the present invention provides a compound of formula (I):



or a salt, solvate and chemically protected form thereof,

10 wherein:

one of R² and R⁵ is:


- (i) H or an optionally substituted C₁₋₄ alkyl group; or
- (ii) an optionally substituted C₅₋₇ aryl; and the other of R² and R⁵ is the other group;

15 m and n can be 0 or 1, and m + n = 1 or 2

R^N is H or optionally substituted C₁₋₄ alkyl


R³ is either:

- (i) carboxy;
- (ii) a group of formula (II):



20

- (iii) a group of formula (III):



wherein R is optionally substituted C₁₋₇ alkyl, C₅₋₂₀ aryl, or NR^{N3}R^{N4}, where R^{N3} and R^{N4} are independently selected from

optionally substituted C₁₋₄ alkyl; or
(iv) tetrazol-5-yl.

A second aspect of the present invention provides a compound
5 of formula (I) or a pharmaceutically acceptable salt thereof
for use in a method of therapy.

A third aspect of the present invention provides a
pharmaceutical composition comprising a compound of formula
10 (I) as defined in the first aspect or a pharmaceutically
acceptable salt thereof together with a pharmaceutically
acceptable carrier or diluent.

A further aspect of the present invention provides the use
15 of a compound of formula (I) or a pharmaceutically
acceptable salt thereof in the preparation of a medicament
for the treatment of a condition alleviated by antagonism of
an EP₄ receptor.

20 Another aspect of the present invention provides a method of
treating a condition which can be alleviated by antagonism
of an EP₄ receptor, which method comprises administering to
a patient in need of treatment an effective amount of a
compound of formula (I), or a pharmaceutically acceptable
25 salt thereof.

Conditions which can be alleviated by antagonism of an EP₄
receptor are discussed above, and particularly include
primary headache disorders, most particularly migraines.

30

The present invention also provides methods of antagonizing
EP₄ receptors, *in vitro* or *in vivo*, comprising contacting a
cell with an effective amount of a compound of formula (I).

In some embodiments, the compounds described above may be selective as against antagonism of the other three EP receptors, i.e. EP₁, EP₂ and EP₃. This selectivity allows for targeting of the effect of the compounds of the invention, with possible benefits in the treatment of certain conditions.

Definitions

Monodentate groups

10 (i.e groups with one point of covalent attachment)

Alkyl: The term "alkyl" as used herein, pertains to a monovalent moiety obtained by removing a hydrogen atom from a carbon atom of a hydrocarbon compound having from 1 to 7 carbon atoms (unless otherwise specified), which may be aliphatic or alicyclic, and which may be saturated or unsaturated. Thus, the term "alkyl" includes the sub-classes alkenyl, alkynyl, cycloalkyl, cycloalkenyl, cylcoalkynyl, etc., discussed below.

20

In the context of alkyl groups, the prefixes (e.g. C₁₋₄, C₁₋₇) denote the number of carbon atoms, or range of number of carbon atoms. For example, the term "C₁₋₄ alkyl" as used herein, pertains to an alkyl group having from 1 to 4 carbon atoms. Examples of groups of alkyl groups include C₁₋₄ alkyl ("lower alkyl") and C₁₋₇ alkyl. Note that the first prefix may vary according to other limitations; for example, for unsaturated alkyl groups, the first prefix must be at least 2; for cyclic alkyl groups, the first prefix must be at least 3; etc.

30

Examples of saturated alkyl groups include, but are not limited to, methyl (C₁), ethyl (C₂), propyl (C₃), butyl (C₄), pentyl (C₅), hexyl (C₆) and heptyl (C₇).

Examples of saturated linear alkyl groups include, but are not limited to, methyl (C_1), ethyl (C_2), n-propyl (C_3), n-butyl (C_4), n-pentyl (amyl) (C_5), n-hexyl (C_6), and n-
5 heptyl (C_7).

Examples of saturated branched alkyl groups include iso-propyl (C_3), iso-butyl (C_4), sec-butyl (C_4), tert-butyl (C_4), iso-pentyl (C_5), and neo-pentyl (C_5).

10

Alkenyl: The term "alkenyl" as used herein, pertains to an alkyl group having one or more carbon-carbon double bonds. Examples of alkenyl groups include C_{2-4} alkenyl and C_{2-7} alkenyl. Examples of alkenyl groups include, but are not
15 limited to, ethenyl (vinyl, $-CH=CH_2$), 1-propenyl ($-CH=CH-CH_3$), 2-propenyl (allyl, $-CH=CH-CH_2$), isopropenyl (1-methylvinyl, $-C(CH_3)=CH_2$), butenyl (C_4), pentenyl (C_5), and hexenyl (C_6).

20 Alkynyl: The term "alkynyl" as used herein, pertains to an alkyl group having one or more carbon-carbon triple bonds. Examples of groups of alkynyl groups include C_{2-4} alkynyl and C_{2-7} alkynyl. Examples of alkynyl groups include, but are not limited to, ethynyl (ethinyl, $-C\equiv CH$) and 2-propynyl
25 (propargyl, $-CH_2-C\equiv CH$).

Cycloalkyl: The term "cycloalkyl" as used herein, pertains to an alkyl group which is also a cyclyl group; that is, a monovalent moiety obtained by removing a hydrogen atom from
30 an alicyclic ring atom of a carbocyclic ring of a carbocyclic compound, which carbocyclic ring may be saturated or unsaturated, which moiety has from 3 to 7 carbon atoms (unless otherwise specified), including from 3 to 7 ring atoms. Thus, the term "cycloalkyl" includes the

sub-classes cycloalkenyl and cycloalkynyl. Preferably, each ring has from 3 to 7 ring atoms. Examples of groups of cycloalkyl groups include C₃₋₇ cycloalkyl.

- 5 Examples of cycloalkyl groups include, but are not limited to, those derived from:

saturated monocyclic hydrocarbon compounds:

- cyclopropane (C₃), cyclobutane (C₄), cyclopentane (C₅),
cyclohexane (C₆), cycloheptane (C₇), methylcyclopropane (C₄),
10 dimethylcyclopropane (C₅), methylcyclobutane (C₅),
dimethylcyclobutane (C₆), methylcyclopentane (C₆),
dimethylcyclopentane (C₇), methylcyclohexane (C₇);

unsaturated monocyclic hydrocarbon compounds:

- cyclopropene (C₃), cyclobutene (C₄), cyclopentene (C₅),
15 cyclohexene (C₆), methylcyclopropene (C₄),
dimethylcyclopropene (C₅), methylcyclobutene (C₅),
dimethylcyclobutene (C₆), methylcyclopentene (C₆),
dimethylcyclopentene (C₇), methylcyclohexene (C₇);

- 20 Heterocyclyl: The term "heterocyclyl" as used herein, pertains to a monovalent moiety obtained by removing a hydrogen atom from a ring atom of a heterocyclic compound, which moiety has from 3 to 20 ring atoms (unless otherwise specified), of which from 1 to 10 are ring heteroatoms.
25 Preferably, each ring has from 3 to 7 ring atoms, of which from 1 to 4 are ring heteroatoms.

- In this context, the prefixes (e.g. C₃₋₂₀, C₃₋₇, C₅₋₆, etc.) denote the number of ring atoms, or range of number of ring
30 atoms, whether carbon atoms or heteroatoms. For example, the term "C₅₋₆ heterocyclyl" as used herein, pertains to a heterocyclyl group having 5 or 6 ring atoms. Examples of groups of heterocyclyl groups include C₃₋₂₀ heterocyclyl, C₅₋₂₀ heterocyclyl, C₃₋₁₅ heterocyclyl, C₅₋₁₅ heterocyclyl, C₃₋₁₂

heterocyclyl, C₅₋₁₂ heterocyclyl, C₃₋₁₀ heterocyclyl, C₅₋₁₀ heterocyclyl, C₃₋₇ heterocyclyl, C₅₋₇ heterocyclyl, and C₅₋₆ heterocyclyl.

- 5 Examples of monocyclic heterocyclyl groups include, but are not limited to, those derived from:

- N₁: aziridine (C₃), azetidine (C₄), pyrrolidine (tetrahydropyrrole) (C₅), pyrroline (e.g., 3-pyrroline, 2,5-dihydropyrrole) (C₅), 2H-pyrrole or 3H-pyrrole
10 (isopyrrole, isoazole) (C₅), piperidine (C₆), dihydropyridine (C₆), tetrahydropyridine (C₆), azepine (C₇);
O₁: oxirane (C₃), oxetane (C₄), oxolane (tetrahydrofuran) (C₅), oxole (dihydrofuran) (C₅), oxane (tetrahydropyran) (C₆), dihydropyran (C₆), pyran (C₆), oxepin (C₇);
15 S₁: thiirane (C₃), thietane (C₄), thiolane (tetrahydrothiophene) (C₅), thiane (tetrahydrothiopyran) (C₆), thiepane (C₇);
O₂: dioxolane (C₅), dioxane (C₆), and dioxepane (C₇);
O₃: trioxane (C₆);
20 N₂: imidazolidine (C₅), pyrazolidine (diazolidine) (C₅), imidazoline (C₅), pyrazoline (dihydropyrazole) (C₅), piperazine (C₆);
N₁O₁: tetrahydrooxazole (C₅), dihydrooxazole (C₅), tetrahydroisoxazole (C₅), dihydroisoxazole (C₅), morpholine (C₆), tetrahydrooxazine (C₆), dihydrooxazine (C₆), oxazine
25 (C₆);
N₁S₁: thiazoline (C₅), thiazolidine (C₅), thiomorpholine (C₆);
N₂O₁: oxadiazine (C₆);
30 O₁S₁: oxathiole (C₅) and oxathiane (thioxane) (C₆); and,
N₁O₁S₁: oxathiazine (C₆).

Aryl: The term "aryl" as used herein, pertains to a monovalent moiety obtained by removing a hydrogen atom from

an aromatic ring atom of an aromatic compound, which moiety has from 3 to 20 ring atoms (unless otherwise specified). Preferably, each ring has from 5 to 7 ring atoms.

- 5 In this context, the prefixes (e.g. C₃₋₂₀, C₅₋₇, C₅₋₆, etc.) denote the number of ring atoms, or range of number of ring atoms, whether carbon atoms or heteroatoms. For example, the term "C₅₋₆ aryl" as used herein, pertains to an aryl group having 5 or 6 ring atoms. Examples of groups of aryl
10 groups include C₃₋₂₀ aryl, C₅₋₂₀ aryl, C₅₋₁₅ aryl, C₅₋₁₂ aryl, C₅₋₁₀ aryl, C₅₋₇ aryl, C₅₋₆ aryl, C₅ aryl, and C₆ aryl.

The ring atoms may be all carbon atoms, as in "carboaryl groups". Examples of carboaryl groups include C₃₋₂₀

- 15 carboaryl, C₅₋₂₀ carboaryl, C₅₋₁₅ carboaryl, C₅₋₁₂ carboaryl, C₅₋₁₀ carboaryl, C₅₋₇ carboaryl, C₅₋₆ carboaryl, C₅ carboaryl, and C₆ carboaryl.

- Examples of carboaryl groups include, but are not limited
20 to, those derived from benzene (i.e. phenyl) (C₆), naphthalene (C₁₀), azulene (C₁₀), anthracene (C₁₄), phenanthrene (C₁₄), naphthacene (C₁₈), and pyrene (C₁₆).

- Examples of aryl groups which comprise fused rings, at least
25 one of which is an aromatic ring, include, but are not limited to, groups derived from indane (e.g., 2,3-dihydro-1H-indene) (C₉), indene (C₉), isoindene (C₉), tetraline (1,2,3,4-tetrahydronaphthalene (C₁₀), acenaphthene (C₁₂), fluorene (C₁₃), phenalene (C₁₃), acephenanthrene (C₁₅), and
30 aceanthrene (C₁₆).

Alternatively, the ring atoms may include one or more heteroatoms, as in "heteroaryl groups". Examples of heteroaryl groups include C₃₋₂₀ heteroaryl, C₅₋₂₀ heteroaryl,

C₅₋₁₅ heteroaryl, C₅₋₁₂ heteroaryl, C₅₋₁₀ heteroaryl, C₅₋₇ heteroaryl, C₅₋₆ heteroaryl, C₅ heteroaryl, and C₆ heteroaryl.

- Examples of monocyclic heteroaryl groups include, but are
5 not limited to, those derived from:
- N₁: pyrrole (azole) (C₅), pyridine (azine) (C₆);
O₁: furan (oxole) (C₅);
S₁: thiophene (thiole) (C₅);
N₁O₁: oxazole (C₅), isoxazole (C₅), isoxazine (C₆);
10 N₂O₁: oxadiazole (furazan) (C₅);
N₃O₁: oxatriazole (C₅);
N₁S₁: thiazole (C₅), isothiazole (C₅);
N₂: imidazole (1,3-diazole) (C₅), pyrazole
(1,2-diazole) (C₅), pyridazine (1,2-diazine) (C₆);
15 pyrimidine (1,3-diazine) (C₆), pyrazine (1,4-diazine) (C₆);
N₃: triazole (C₅), triazine (C₆); and,
N₄: tetrazole (C₅).

- Examples of heteroaryl groups which comprise fused rings,
20 include, but are not limited to:
- C₉ (with 2 fused rings) derived from benzofuran (O₁),
isobenzofuran (O₁), indole (N₁), isoindole (N₁), indolizine
(N₁), indoline (N₁), isoindoline (N₁), purine (N₄) (e.g.,
adenine, guanine), benzimidazole (N₂), indazole (N₂),
25 benzoxazole (N₁O₁), benzisoxazole (N₁O₁), benzodioxole (O₂),
benzofurazan (N₂O₁), benzotriazole (N₃), benzothiofuran (S₁),
benzothiazole (N₁S₁), benzothiadiaazole (N₂S);
- C₁₀ (with 2 fused rings) derived from chromene (O₁),
isochromene (O₁), chroman (O₁), isochroman (O₁), benzodioxan
30 (O₂), quinoline (N₁), isoquinoline (N₁), quinolizine (N₁),
benzoxazine (N₁O₁), benzodiazine (N₂), pyridopyridine (N₂),
quinoxaline (N₂), quinazoline (N₂), cinnoline (N₂),
phthalazine (N₂), naphthyridine (N₂), pteridine (N₄);

C₁₁ (with 2 fused rings) derived from benzodiazepine (N₂);

C₁₃ (with 3 fused rings) derived from carbazole (N₁), dibenzofuran (O₁), dibenzothiophene (S₁), carboline (N₂),
5 perimidine (N₂), pyridoindole (N₂); and,

C₁₄ (with 3 fused rings) derived from acridine (N₁), xanthene (O₁), thioxanthene (S₁), oxanthrene (O₂), phenoxathiin (O₁S₁), phenazine (N₂), phenoxazine (N₁O₁), phenothiazine (N₁S₁), thianthrene (S₂), phenanthridine (N₁),
10 phenanthroline (N₂), phenazine (N₂).

If a heteroaryl or heterocyclyl group contains a nitrogen ring atom, this ring atom, where possible, may be in a oxidised state, as an N-oxide.

15

The above groups, whether alone or part of another substituent, may themselves optionally be substituted with one or more groups selected from themselves, the additional monodentate substituents listed below and alkoxylenes.

20

Halo: -F, -Cl, -Br, and -I.

Hydroxy: -OH.

25 Ether: -OR, wherein R is an ether substituent, for example, a C₁₋₇ alkyl group (also referred to as a C₁₋₇ alkoxy group, discussed below), a C₃₋₂₀ heterocyclyl group (also referred to as a C₃₋₂₀ heterocycliloxy group), or a C₅₋₂₀ aryl group (also referred to as a C₅₋₂₀ aryloxy group), preferably a C₁₋₇
30 alkyl group.

C₁₋₇ alkoxy: -OR, wherein R is a C₁₋₇ alkyl group. Examples of C₁₋₇ alkoxy groups include, but are not limited to, -OMe (methoxy), -OEt (ethoxy), -O(nPr) (n-propoxy), -O(iPr)

(isopropoxy), -O(nBu) (n-butoxy), -O(sBu) (sec-butoxy),
-O(iBu) (isobutoxy), and -O(tBu) (tert-butoxy).

Oxo (keto, -one): =O.

5

Thione (thioketone): =S.

Imino (imine): =NR, wherein R is an imino substituent, for
example, hydrogen, C₁₋₇ alkyl group, a C₃₋₂₀ heterocyclyl
10 group, or a C₅₋₂₀ aryl group, preferably hydrogen or a C₁₋₇
alkyl group. Examples of imino groups include, but are not
limited to, =NH, =NMe, =NEt, and =NPh.

Formyl (carbaldehyde, carboxaldehyde): -C(=O)H.

15

Acyl (keto): -C(=O)R, wherein R is an acyl substituent, for
example, a C₁₋₇ alkyl group (also referred to as C₁₋₇
alkylacyl or C₁₋₇ alkanoyl), a C₃₋₂₀ heterocyclyl group (also
referred to as C₃₋₂₀ heterocyclylacyl), or a C₅₋₂₀ aryl group
20 (also referred to as C₅₋₂₀ arylacyl), preferably a C₁₋₇ alkyl
group. Examples of acyl groups include, but are not limited
to, -C(=O)CH₃ (acetyl), -C(=O)CH₂CH₃ (propionyl),
-C(=O)C(CH₃)₃ (t-butyryl), and -C(=O)Ph (benzoyl, phenone).

25 Carboxy (carboxylic acid): -C(=O)OH.

Thiocarboxy (thiocarboxylic acid): -C(=S)SH.

Thiolocarboxy (thiolocarboxylic acid): -C(=O)SH.

30

Thionocarboxy (thionocarboxylic acid): -C(=S)OH.

Imidic acid: -C(=NH)OH.

Hydroxamic acid: $-C(=NOH)OH$.

Ester (carboxylate, carboxylic acid ester, oxycarbonyl):

$-C(=O)OR$, wherein R is an ester substituent, for example, a
5 C_{1-7} alkyl group, a C_{3-20} heterocyclyl group, or a C_{5-20} aryl
group, preferably a C_{1-7} alkyl group. Examples of ester
groups include, but are not limited to, $-C(=O)OCH_3$,
 $-C(=O)OCH_2CH_3$, $-C(=O)OC(CH_3)_3$, and $-C(=O)OPh$.


- 10 Acyloxy (reverse ester): $-OC(=O)R$, wherein R is an acyloxy
substituent, for example, a C_{1-7} alkyl group, a C_{3-20}
heterocyclyl group, or a C_{5-20} aryl group, preferably a C_{1-7}
alkyl group. Examples of acyloxy groups include, but are
not limited to, $-OC(=O)CH_3$ (acetoxy), $-OC(=O)CH_2CH_3$,
15 $-OC(=O)C(CH_3)_3$, $-OC(=O)Ph$, and $-OC(=O)CH_2Ph$.

Amido (carbamoyl, carbamyl, aminocarbonyl, carboxamide):

- $-C(=O)NR^1R^2$, wherein R^1 and R^2 are independently amino
substituents, as defined for amino groups. Examples of
20 amido groups include, but are not limited to, $-C(=O)NH_2$,
 $-C(=O)NHCH_3$, $-C(=O)N(CH_3)_2$, $-C(=O)NHCH_2CH_3$, and
 $-C(=O)N(CH_2CH_3)_2$, as well as amido groups in which R^1 and R^2 ,
together with the nitrogen atom to which they are attached,
form a heterocyclic structure as in, for example,
25 piperidinocarbonyl, morpholinocarbonyl,
thiomorpholinocarbonyl, and piperazinocarbonyl.

- Acylamino: $-NR^1C(=O)R^2$, wherein R^1 is an amide substituent,
for example, hydrogen, a C_{1-7} alkyl group, a C_{3-20} heterocyclyl
30 group, or a C_{5-20} aryl group, preferably hydrogen or a C_{1-7}
alkyl group, and R^2 is an acyl substituent, for example, a
 C_{1-7} alkyl group, a C_{3-20} heterocyclyl group, or a C_{5-20} aryl
group, preferably hydrogen or a C_{1-7} alkyl group. Examples
of acylamide groups include, but are not limited to,

-NHC(=O)CH_3 , $\text{-NHC(=O)CH}_2\text{CH}_3$, and -NHC(=O)Ph . R^1 and R^2 may together form a cyclic structure, as in, for example, succinimidyl, maleimidyl, and phthalimidyl:



5

Thioamido (thiocarbamyl): $\text{-C(=S)NR}^1\text{R}^2$, wherein R^1 and R^2 are independently amino substituents, as defined for amino groups. Examples of thioamido groups include, but are not limited to, -C(=S)NH_2 , -C(=S)NHCH_3 , $\text{-C(=S)N(CH}_3)_2$, and


10 $\text{-C(=S)NHCH}_2\text{CH}_3$.

Ureido: $\text{-N(R}^1\text{)CONR}^2\text{R}^3$ wherein R^2 and R^3 are independently amino substituents, as defined for amino groups, and R^1 is a ureido substituent, for example, hydrogen, a C_{1-7} alkyl group, a C_{3-20} heterocyclyl group, or a C_{5-20} aryl group, preferably hydrogen or a C_{1-7} alkyl group. Examples of ureido groups include, but are not limited to, -NHCONH_2 , -NHCONHMe , -NHCONHEt , -NHCONMe_2 , -NHCONEt_2 , -NMeCONH_2 , -NMeCONHMe , -NMeCONHEt , -NMeCONMe_2 , and -NMeCONEt_2 .

20

Guanidino: -NH-C(=NH)NH_2 .

Tetrazolyl: a five membered aromatic ring having four nitrogen atoms and one carbon atom,



25

Amino: $-NR^1R^2$, wherein R^1 and R^2 are independently amino substituents, for example, hydrogen, a C_{1-7} alkyl group (also referred to as C_{1-7} alkylamino or di- C_{1-7} alkylamino), a C_{3-20} heterocyclyl group, or a C_{5-20} aryl group, preferably H or a C_{1-7} alkyl group, or, in the case of a "cyclic" amino group, R^1 and R^2 , taken together with the nitrogen atom to which they are attached, form a heterocyclic ring having from 4 to 8 ring atoms. Amino groups may be primary ($-NH_2$), secondary ($-NHR^1$), or tertiary ($-NHR^1R^2$), and in cationic form, may be quaternary ($-^+NR^1R^2R^3$). Examples of amino groups include, but are not limited to, $-NH_2$, $-NHCH_3$, $-NHC(CH_3)_2$, $-N(CH_3)_2$, $-N(CH_2CH_3)_2$, and $-NHPh$. Examples of cyclic amino groups include, but are not limited to, aziridino, azetidino, pyrrolidino, piperidino, piperazino, morpholino, and thiomorpholino.

Amidine (amidino): $-C(=NR)NR_2$, wherein each R is an amidine substituent, for example, hydrogen, a C_{1-7} alkyl group, a C_{3-20} heterocyclyl group, or a C_{5-20} aryl group, preferably H or a C_{1-7} alkyl group. Examples of amidine groups include, but are not limited to, $-C(=NH)NH_2$, $-C(=NH)NMe_2$, and $-C(=NMe)NMe_2$.

Nitro: $-NO_2$.

Nitroso: $-NO$.

Cyano (nitrile, carbonitrile): $-CN$.

Sulfhydryl (thiol, mercapto): $-SH$.

Thioether (sulfide): $-SR$, wherein R is a thioether substituent, for example, a C_{1-7} alkyl group (also referred to as a C_{1-7} alkylthio group), a C_{3-20} heterocyclyl group, or a

C₅₋₂₀ aryl group, preferably a C₁₋₇ alkyl group. Examples of C₁₋₇ alkylthio groups include, but are not limited to, -SCH₃ and -SCH₂CH₃.

- 5 Disulfide: -SS-R, wherein R is a disulfide substituent, for example, a C₁₋₇ alkyl group, a C₃₋₂₀ heterocyclyl group, or a C₅₋₂₀ aryl group, preferably a C₁₋₇ alkyl group (also referred to herein as C₁₋₇ alkyl disulfide). Examples of C₁₋₇ alkyl disulfide groups include, but are not limited to, -SSCH₃ and
10 -SSCH₂CH₃.

- Sulfine (sulfinyl, sulfoxide): -S(=O)R, wherein R is a sulfine substituent, for example, a C₁₋₇ alkyl group, a C₃₋₂₀ heterocyclyl group, or a C₅₋₂₀ aryl group, preferably a C₁₋₇
15 alkyl group. Examples of sulfine groups include, but are not limited to, -S(=O)CH₃ and -S(=O)CH₂CH₃.

- Sulfone (sulfonyl): -S(=O)₂R, wherein R is a sulfone substituent, for example, a C₁₋₇ alkyl group, a C₃₋₂₀
20 heterocyclyl group, or a C₅₋₂₀ aryl group, preferably a C₁₋₇ alkyl group, including, for example, a fluorinated or perfluorinated C₁₋₇ alkyl group. Examples of sulfone groups include, but are not limited to, -S(=O)₂CH₃ (methanesulfonyl, mesyl), -S(=O)₂CF₃ (triflyl), -S(=O)₂CH₂CH₃ (esyl), -S(=O)₂C₄F₉ (nonafllyl), -S(=O)₂CH₂CF₃ (tresyl),
25 -S(=O)₂CH₂CH₂NH₂ (tauryl), -S(=O)₂Ph (phenylsulfonyl, besyl), 4-methylphenylsulfonyl (tosyl), 4-chlorophenylsulfonyl (closyl), 4-bromophenylsulfonyl (brosyl), 4-nitrophenyl (nosyl), 2-naphthalenesulfonate (napsyl), and
30 5-dimethylamino-naphthalen-1-ylsulfonate (dansyl).

Sulfinic acid (sulfino): -S(=O)OH, -SO₂H.

Sulfonic acid (sulfo): -S(=O)₂OH, -SO₃H.

Sulfinate (sulfinic acid ester): $-S(=O)OR$; wherein R is a sulfinate substituent, for example, a C_{1-7} alkyl group, a C_{3-20} heterocyclyl group, or a C_{5-20} aryl group, preferably a C_{1-7} alkyl group. Examples of sulfinate groups include, but are not limited to, $-S(=O)OCH_3$ (methoxysulfinyl; methyl sulfinate) and $-S(=O)OCH_2CH_3$ (ethoxysulfinyl; ethyl sulfinate).

10 Sulfinyloxy: $-OS(=O)R$, wherein R is a sulfinyloxy substituent, for example, a C_{1-7} alkyl group, a C_{3-20} heterocyclyl group, or a C_{5-20} aryl group, preferably a C_{1-7} alkyl group. Examples of sulfinyloxy groups include, but are not limited to, $-OS(=O)CH_3$ and $-OS(=O)CH_2CH_3$.

15

Sulfamyl (sulfamoyl; sulfinic acid amide; sulfinamide): $-S(=O)NR^1R^2$, wherein R^1 and R^2 are independently amino substituents, as defined for amino groups. Examples of sulfamyl groups include, but are not limited to, $-S(=O)NH_2$, $-S(=O)NH(CH_3)$, $-S(=O)N(CH_3)_2$, $-S(=O)NH(CH_2CH_3)$, $-S(=O)N(CH_2CH_3)_2$, and $-S(=O)NHPh$.

20

Sulfonamido (sulfinamoyl; sulfonic acid amide; sulfonamide): $-S(=O)_2NR^1R^2$, wherein R^1 and R^2 are independently amino substituents, as defined for amino groups. Examples of sulfonamido groups include, but are not limited to, $-S(=O)_2NH_2$, $-S(=O)_2NH(CH_3)$, $-S(=O)_2N(CH_3)_2$, $-S(=O)_2NH(CH_2CH_3)$, $-S(=O)_2N(CH_2CH_3)_2$, and $-S(=O)_2NHPh$.

25

30 Sulfonamino: $-NR^1S(=O)_2R$, wherein R^1 is an amino substituent, as defined for amino groups, and R is a sulfonamino substituent, for example, a C_{1-7} alkyl group, a C_{3-20} heterocyclyl group, or a C_{5-20} aryl group, preferably a C_{1-7}

alkyl group. Examples of sulfonamino groups include, but are not limited to, $\text{-NHS(=O)}_2\text{CH}_3$ and $\text{-N(CH}_3\text{)S(=O)}_2\text{C}_6\text{H}_5$.

5 Sulfinamino: $\text{-NR}^1\text{S(=O)R}$, wherein R^1 is an amino substituent, as defined for amino groups, and R is a sulfinamino substituent, for example, a C_{1-7} alkyl group, a C_{3-20} heterocyclyl group, or a C_{5-20} aryl group, preferably a C_{1-7} alkyl group. Examples of sulfinamino groups include, but are not limited to, -NHS(=O)CH_3 and $\text{-N(CH}_3\text{)S(=O)C}_6\text{H}_5$.

10

As already mentioned, the above described groups may be substituted, and particular examples include, but are not limited to, C_{3-20} aryl- C_{1-7} alkyl groups, which include benzyl (phenylmethyl, $\text{PhCH}_2\text{-}$), benzhydryl ($\text{Ph}_2\text{CH-}$), trityl (triphenylmethyl, $\text{Ph}_3\text{C-}$), phenethyl (phenylethyl, $\text{Ph-CH}_2\text{CH}_2\text{-}$), styryl (Ph-CH=CH-) and cinnamyl ($\text{Ph-CH=CH-CH}_2\text{-}$).

15

Bidentate groups

(i.e. groups with two points of covalent attachment; linking groups)

20

Alkylene: The term " C_{1-3} alkylene", as used herein, pertains to a bidentate moiety obtained by removing two hydrogen atoms from each of two different carbon atoms, of a linear hydrocarbon compound having from 1 to 3 carbon atoms, which may be saturated or unsaturated. Thus, the term "alkylene" includes the sub-classes alkenylene and alkynylene.

25

In this context, the prefix C_{1-3} denotes the number of carbon atoms, or range of number of carbon atoms.

30

Examples of saturated C_{1-3} alkylene groups include $\text{-CH}_2\text{-}$ (methylene), $\text{-CH}_2\text{CH}_2\text{-}$ (ethylene) and $\text{-CH}_2\text{CH}_2\text{CH}_2\text{-}$ (propylene).

Examples of unsaturated C₁₋₃ alkylene groups (which may be termed "C₂₋₃ alkenylene" or "C₂₋₃ alkynylene", as appropriate) include -CH=CH- (vinylene), -CH=CH-CH₂-, -CH₂-CH=CH-, -C≡C-, -C≡C-CH₂- and -CH₂-C≡C-.

5

The C₁₋₃ alkylene group may be substituted by any monodentate substituent described above.

10 Alkoxylenes: The term "alkoxylenes," as used herein, pertains to a bidentate group of formula -O(CH₂)_nO-, where n is 1 or 2.

Includes Other Forms

15 Unless otherwise specified, included in the above are the well known ionic, salt, solvate, and protected forms of these substituents. For example, a reference to carboxylic acid (-COOH) also includes the anionic (carboxylate) form (-COO⁻), a salt or solvate thereof, as well as conventional protected forms. Similarly, a reference to an amino group 20 includes the protonated form (-N⁺HR¹R²), a salt or solvate of the amino group, for example, a hydrochloride salt, as well as conventional protected forms of an amino group. Similarly, a reference to a hydroxyl group also includes the anionic form (-O⁻), a salt or solvate thereof, as well as 25 conventional protected forms of a hydroxyl group.


Isomers, Salts, Solvates and Protected Forms

30 Certain compounds may exist in one or more particular geometric, optical, enantiomeric, diastereomeric, epimeric, stereoisomeric, tautomeric, conformational, or anomeric forms, including but not limited to, cis- and trans-forms; E- and Z-forms; c-, t-, and r- forms; endo- and exo-forms; R-, S-, and meso-forms; D- and L-forms; d- and l-forms; (+) and (-) forms; keto-, enol-, and enolate-forms; syn- and

anti-forms; synclinal- and anticlinal-forms; α - and β -forms; axial and equatorial forms; boat-, chair-, twist-, envelope-, and halfchair-forms; and combinations thereof, hereinafter collectively referred to as "isomers" (or
5 "isomeric forms").

Note that, except as discussed below for tautomeric forms, specifically excluded from the term "isomers", as used herein, are structural (or constitutional) isomers
10 (i.e. isomers which differ in the connections between atoms rather than merely by the position of atoms in space). For example, a reference to a methoxy group, $-\text{OCH}_3$, is not to be construed as a reference to its structural isomer, a hydroxymethyl group, $-\text{CH}_2\text{OH}$. Similarly, a reference to
15 ortho-chlorophenyl is not to be construed as a reference to its structural isomer, meta-chlorophenyl. However, a reference to a class of structures may well include structurally isomeric forms falling within that class (e.g. C_{1-7} alkyl includes n-propyl and iso-propyl; butyl
20 includes n-, iso-, sec-, and tert-butyl; methoxyphenyl includes ortho-, meta-, and para-methoxyphenyl).

The above exclusion does not pertain to tautomeric forms, for example, keto-, enol-, and enolate-forms, as in, for
25 example, the following tautomeric pairs: keto/enol (illustrated below), imine/enamine, amide/imino alcohol, amidine/amidine, nitroso/oxime, thioketone/enethiol, N-nitroso/hydroxyazo, and nitro/aci-nitro.



30

Note that specifically included in the term "isomer" are compounds with one or more isotopic substitutions. For

example, H may be in any isotopic form, including ^1H , ^2H (D), and ^3H (T); C may be in any isotopic form, including ^{12}C , ^{13}C , and ^{14}C ; O may be in any isotopic form, including ^{16}O and ^{18}O ; and the like.

5

Unless otherwise specified, a reference to a particular compound includes all such isomeric forms, including (wholly or partially) racemic and other mixtures thereof. Methods for the preparation (e.g. asymmetric synthesis) and
10 separation (e.g. fractional crystallisation and chromatographic means) of such isomeric forms are either known in the art or are readily obtained by adapting the methods taught herein, or known methods, in a known manner.

15 Unless otherwise specified, a reference to a particular compound also includes ionic, salt, solvate, and protected forms of thereof, for example, as discussed below.

It may be convenient or desirable to prepare, purify, and/or
20 handle a corresponding salt of the active compound, for example, a pharmaceutically-acceptable salt. Examples of pharmaceutically acceptable salts are discussed in Berge, et al., *J. Pharm. Sci.*, **66**, 1-19 (1977).

25 For example, if the compound is anionic, or has a functional group which may be anionic (e.g. $-\text{COOH}$ may be $-\text{COO}^-$), then a salt may be formed with a suitable cation. Examples of suitable inorganic cations include, but are not limited to, alkali metal ions such as Na^+ and K^+ , alkaline earth cations
30 such as Ca^{2+} and Mg^{2+} , and other cations such as Al^{+3} . Examples of suitable organic cations include, but are not limited to, ammonium ion (i.e. NH_4^+) and substituted ammonium ions (e.g. NH_3R^+ , NH_2R_2^+ , NHR_3^+ , NR_4^+). Examples of some suitable substituted ammonium ions are those derived

from: ethylamine, diethylamine, dicyclohexylamine, triethylamine, butylamine, ethylenediamine, ethanolamine, diethanolamine, piperazine, benzylamine, phenylbenzylamine, choline, meglumine, and tromethamine, as well as amino
5 acids, such as lysine and arginine. An example of a common quaternary ammonium ion is $N(CH_3)_4^+$.

If the compound is cationic, or has a functional group which may be cationic (e.g. $-NH_2$ may be $-NH_3^+$), then a salt may be
10 formed with a suitable anion. Examples of suitable inorganic anions include, but are not limited to, those derived from the following inorganic acids: hydrochloric, hydrobromic, hydroiodic, sulfuric, sulfurous, nitric, nitrous, phosphoric, and phosphorous.

15 Examples of suitable organic anions include, but are not limited to, those derived from the following organic acids: 2-acetyoxybenzoic, acetic, ascorbic, aspartic, benzoic, camphorsulfonic, cinnamic, citric, edetic, ethanedisulfonic,
20 ethanesulfonic, fumaric, glucoheptonic, gluconic, glutamic, glycolic, hydroxymaleic, hydroxynaphthalene carboxylic, isethionic, lactic, lactobionic, lauric, maleic, malic, methanesulfonic, mucic, oleic, oxalic, palmitic, pantoic, pantothenic, phenylacetic, phenylsulfonic, propionic,
25 pyruvic, salicylic, stearic, succinic, sulfanilic, tartaric, toluenesulfonic, and valeric. Examples of suitable polymeric organic anions include, but are not limited to, those derived from the following polymeric acids: tannic acid, carboxymethyl cellulose.

30

It may be convenient or desirable to prepare, purify, and/or handle a corresponding solvate of the active compound. The term "solvate" is used herein in the conventional sense to refer to a complex of solute (e.g., active compound, salt of

active compound) and solvent. If the solvent is water, the solvate may be conveniently referred to as a hydrate, for example, a mono-hydrate, a di-hydrate, a tri-hydrate, etc.

5 It may be convenient or desirable to prepare, purify, and/or handle the active compound in a chemically protected form. The term "chemically protected form" is used herein in the conventional chemical sense and pertains to a compound in which one or more reactive functional groups are protected
10 from undesirable chemical reactions under specified conditions (e.g. pH, temperature, radiation, solvent, and the like). In practice, well known chemical methods are employed to reversibly render unreactive a functional group, which otherwise would be reactive, under specified
15 conditions. In a chemically protected form, one or more reactive functional groups are in the form of a protected or protecting group (also known as a masked or masking group or a blocked or blocking group). By protecting a reactive functional group, reactions involving other unprotected
20 reactive functional groups can be performed, without affecting the protected group; the protecting group may be removed, usually in a subsequent step, without substantially affecting the remainder of the molecule. See, for example, Protective Groups in Organic Synthesis (T. Green and
25 P. Wuts; 3rd Edition; John Wiley and Sons, 1999).

A wide variety of such "protecting", "blocking", or "masking" methods are widely used and well known in organic synthesis. For example, a compound which has two
30 nonequivalent reactive functional groups, both of which would be reactive under specified conditions, may be derivatized to render one of the functional groups "protected," and therefore unreactive, under the specified conditions; so protected, the compound may be used as a

reactant which has effectively only one reactive functional group. After the desired reaction (involving the other functional group) is complete, the protected group may be "deprotected" to return it to its original functionality.

5

For example, a hydroxy group may be protected as an ether (-OR) or an ester (-OC(=O)R), for example, as: a t-butyl ether; a benzyl, benzhydryl (diphenylmethyl), or trityl (triphenylmethyl) ether; a trimethylsilyl or
10 t-butyldimethylsilyl ether; or an acetyl ester (-OC(=O)CH₃, -OAc).

For example, an aldehyde or ketone group may be protected as an acetal (R-CH(OR)₂) or ketal (R₂C(OR)₂), respectively, in
15 which the carbonyl group (>C=O) is converted to a diether (>C(OR)₂), by reaction with, for example, a primary alcohol.

The aldehyde or ketone group is readily regenerated by hydrolysis using a large excess of water in the presence of acid.

20

For example, an amine group may be protected, for example, as an amide (-NRCO-R) or a urethane (-NRCO-OR), for example, as: an acetamide (-NHCO-CH₃); a benzyloxy amide (-NHCO-OCH₂C₆H₅, -NH-Cbz); as a t-butoxy amide (-NHCO-OC(CH₃)₃, -NH-Boc); a 2-biphenyl-2-propoxy amide (-NHCO-OC(CH₃)₂C₆H₄C₆H₅, -NH-Bpoc), as a 9-fluorenylmethoxy amide (-NH-Fmoc), as a 6-nitroveratryloxy amide (-NH-Nvoc), as a 2-trimethylsilylethyloxy amide (-NH-Teoc), as a 2,2,2-trichloroethyloxy amide (-NH-Troc), as an allyloxy amide
30 (-NH-Alloc), as a 2(-phenylsulfonyl)ethyloxy amide (-NH-Psec); or, in suitable cases (e.g., cyclic amines), as a nitroxide radical (>N-O·).

For example, a carboxylic acid group may be protected as an ester for example, as: an C₁₋₇ alkyl ester (e.g., a methyl ester; a t-butyl ester); a C₁₋₇ haloalkyl ester (e.g., a C₁₋₇ trihaloalkyl ester); a triC₁₋₇ alkylsilyl-C₁₋₇ alkyl ester; or
5 a C₅₋₂₀ aryl-C₁₋₇ alkyl ester (e.g. a benzyl ester; a nitrobenzyl ester); or as an amide, for example, as a methyl amide.

For example, a thiol group may be protected as a thioether
10 (-SR), for example, as: a benzyl thioether; an acetamidomethyl ether (-S-CH₂NHC(=O)CH₃).

The term "treatment", as used herein in the context of treating a condition, pertains generally to treatment and
15 therapy, whether of a human or an animal (e.g. in veterinary applications), in which some desired therapeutic effect is achieved, for example, the inhibition of the progress of the condition, and includes a reduction in the rate of progress, a halt in the rate of progress, amelioration of the
20 condition, and cure of the condition. Treatment as a prophylactic measure (i.e. prophylaxis) is also included.

The term "therapeutically-effective amount", as used herein, pertains to that amount of an active compound, or a
25 material, composition or dosage form comprising an active compound, which is effective for producing some desired therapeutic effect, commensurate with a reasonable benefit/risk ratio, when administered in accordance with a desired treatment regimen. Suitable dose ranges will
30 typically be in the range of from 0.01 to 20 mg/kg/day, preferably from 0.1 to 10 mg/kg/day.

Compositions and their administration

Compositions may be formulated for any suitable route and

means of administration. Pharmaceutically acceptable carriers or diluents include those used in formulations suitable for oral, rectal, nasal, topical (including buccal and sublingual), vaginal or parenteral (including

5 subcutaneous, intramuscular, intravenous, intradermal, intrathecal and epidural) administration. The formulations may conveniently be presented in unit dosage form and may be prepared by any of the methods well known in the art of pharmacy. Such methods include the step of bringing into

10 association the active ingredient with the carrier which constitutes one or more accessory ingredients. In general the formulations are prepared by uniformly and intimately bringing into association the active ingredient with liquid carriers or finely divided solid carriers or both, and then,

15 if necessary, shaping the product.

For solid compositions, conventional non-toxic solid carriers include, for example, pharmaceutical grades of mannitol, lactose, cellulose, cellulose derivatives, starch,

20 magnesium stearate, sodium saccharin, talcum, glucose, sucrose, magnesium carbonate, and the like may be used. The active compound as defined above may be formulated as suppositories using, for example, polyalkylene glycols, acetylated triglycerides and the like, as the carrier.

25 Liquid pharmaceutically administrable compositions can, for example, be prepared by dissolving, dispersing, etc, an active compound as defined above and optional pharmaceutical adjuvants in a carrier, such as, for example, water, saline aqueous dextrose, glycerol, ethanol, and the like, to

30 thereby form a solution or suspension. If desired, the pharmaceutical composition to be administered may also contain minor amounts of non-toxic auxiliary substances such as wetting or emulsifying agents, pH buffering agents and the like, for example, sodium acetate, sorbitan monolaurate,

triethanolamine sodium acetate, sorbitan monolaurate, triethanolamine oleate, etc. Actual methods of preparing such dosage forms are known, or will be apparent, to those skilled in this art; for example, see Remington's
5 Pharmaceutical Sciences, 20th edition, pub. Lippincott, Williams & Wilkins, 2000. The composition or formulation to be administered will, in any event, contain a quantity of the active compound(s) in an amount effective to alleviate the symptoms of the subject being treated.

10 Dosage forms or compositions containing active ingredient in the range of 0.25 to 95% with the balance made up from non-toxic carrier may be prepared.

15 For oral administration, a pharmaceutically acceptable non-toxic composition is formed by the incorporation of any of the normally employed excipients, such as, for example, pharmaceutical grades of mannitol, lactose, cellulose, cellulose derivatives, sodium crosscarmellose, starch,
20 magnesium stearate, sodium saccharin, talcum, glucose, sucrose, magnesium carbonate, and the like. Such compositions take the form of solutions, suspensions, tablets, pills, capsules, powders, sustained release formulations and the like. Such compositions may contain
25 1%-95% active ingredient, more preferably 2-50%, most preferably 5-8%.

Parenteral administration is generally characterized by injection, either subcutaneously, intramuscularly or
30 intravenously. Injectables can be prepared in conventional forms, either as liquid solutions or suspensions, solid forms suitable for solution or suspension in liquid prior to injection, or as emulsions. Suitable excipients are, for example, water, saline, dextrose, glycerol, ethanol or the

like. In addition, if desired, the pharmaceutical compositions to be administered may also contain minor amounts of non-toxic auxiliary substances such as wetting or emulsifying agents, pH buffering agents and the like, such
5 as for example, sodium acetate, sorbitan monolaurate, triethanolamine oleate, triethanolamine sodium acetate, etc.

The percentage of active compound contained in such parental compositions is highly dependent on the specific nature
10 thereof, as well as the activity of the compound and the needs of the subject. However, percentages of active ingredient of 0.1% to 10% in solution are employable, and will be higher if the composition is a solid which will be subsequently diluted to the above percentages. Preferably,
15 the composition will comprise 0.2-2% of the active agent in solution.


Acronyms

For convenience, many chemical moieties are represented
20 using well known abbreviations, including but not limited to, methyl (Me), ethyl (Et), n-propyl (nPr), iso-propyl (iPr), n-butyl (nBu), sec-butyl (sBu), iso-butyl (iBu), tert-butyl (tBu), n-hexyl (nHex), cyclohexyl (cHex), phenyl (Ph), biphenyl (biPh), benzyl (Bn), naphthyl (naph), methoxy
25 (MeO), ethoxy (EtO), benzoyl (Bz), and acetyl (Ac).


For convenience, many chemical compounds are represented using well known abbreviations, including but not limited to, methanol (MeOH), ethanol (EtOH), iso-propanol (i-PrOH),
30 methyl ethyl ketone (MEK), ether or diethyl ether (Et₂O), acetic acid (AcOH), dichloromethane (methylene chloride, DCM), acetonitrile (ACN), trifluoroacetic acid (TFA), dimethylformamide (DMF), tetrahydrofuran (THF), and dimethylsulfoxide (DMSO).

General Synthesis Methods

Compounds of the invention wherein R³ is of formula (II):




- 5 may be synthesised from the analogous compound of the invention wherein R³ is carboxy, by reaction with a compound of formula 1:




- 10 in basic conditions, preferably aided by a coupling agent, for example, 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride.

Compounds of the invention wherein R³ is of formula (III):



- 15 may be synthesized from a compound of formula 2:




by reaction with a compound of formula 3:




- 20 wherein X is either OH or halo, where if X is OH, the use of basic conditions and a coupling agent is preferred.

Compounds where R^3 is tetrazol-5-yl may be synthesised from compounds of formula 4:




Formula 4

- 5 by treatment with sodium azide in the presence of a base. Compounds of formula 4 may be synthesised by coupling compounds of Formula 5 and Formula 6a.




Formula 5



Formula 6a

- 10 Such a coupling step may be carried out using a coupling agent, for example, O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate.


Compounds where R^3 is carboxy, may be synthesised from compounds of formula 7:




Formula 7

, where R^0 is typically a C_{1-4} alkyl group, by a hydrolysis reaction, for example, using sodium hydroxide.

Compounds of formula 7 can be synthesised by coupling
5 compounds of formula 5 and 6b:




Formula 5



Formula 6b



Such a coupling step may be carried out as described above,
by using a coupling agent, for example, O-(7-
azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium
10 hexafluorophosphate.

In compounds of formula 5, if R^5 is an aryl group, then
these may be synthesised from compounds of formula 8:



Formula 8


15 by a Suzuki coupling of a compound of formula 9a (or
equivalent ester of formula 9b):



Formula 9b

The Suzuki coupling may be achieved using, for example, [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium (II) as the palladium catalyst.

- 5 Compounds of Formula 8 may be synthesised from compounds of formula 10:



Formula 10

by treating the compound of formula 10 with a brominating agent, such as pyridinium tribromide.

10

Compounds where R² or R⁵ is a phenyl group substituted by -O-CHF₂, can be synthesised from the corresponding compound where the phenyl group is substituted by -OH, by treating this compound with a base and chlorodifluoromethane.

15

Preferences

The following preferences may be combined with one another, and may be different for each aspect of the present invention.

20

R⁵ is preferably the optionally substituted C₅₋₇ aryl group and R² is preferably H or the optionally substituted C₁₋₄ alkyl group.

25

R² is preferably selected from H or an optionally substituted C₁₋₃ alkyl group, more preferably H, methyl, CF₃ or iso-propyl, and most preferably R² is a methyl group.


30

R⁵ is preferably a C₆ aryl group, and is more preferably phenyl. R⁵ may be substituted, and preferred substituents include C₁₋₇ alkoxy groups, more preferably C₁₋₄ alkoxy

groups, e.g. -OMe, -OCF₃, -OEt, -OCHF₂, with -OCHF₂ being the most preferred.


R³ is preferably either:

- 5 (i) a group of formula (II):



; or

- (ii) a group of formula (III):



- 10 with a group of formula (II) being more preferred. In some embodiments, R³ is preferably carboxy.

Where R³ is of formula (II) or (III), R is preferably selected from an optionally substituted C₅₋₂₀ aryl group, and an optionally substituted C₅₋₂₀ aryl-C₁₋₇ alkyl group, wherein
15 the C₁₋₇ alkyl group is more preferably methyl. In these groups the C₅₋₂₀ aryl group is preferably a C₆ aryl group. Such groups may preferably be substituted with C₁₋₄ alkyl groups, such as methyl and hydroxy or halo groups, for example, fluoro. Thus, preferred R groups include, but are
20 not limited to: phenyl; benzyl; 2-fluoro-phenyl; 4-hydroxy-phenyl; 2-trifluoromethyl-phenyl; 5-methyl-pyrid-2-yl.

If R in formula (II) or (III) is a C₁₋₇ alkyl group, it is more preferably a C₁₋₄ alkyl group, for example methyl or
25 propyl.

Preferably n + m = 1, and more preferably n is 0 and m is 1.

R^N is preferably H or methyl, and is more preferably H.

Particularly preferred compounds of the present invention include:

- 5 (4-{[5-(4-Methoxy-phenyl)-2-trifluoromethyl-furan-3-carbonyl]-amino}-phenyl)-acetic acid (4);
(4-{[5-(4-Methoxy-phenyl)-2-methyl-furan-3-carbonyl]-amino}-phenyl)-acetic acid (6);
(4-{[5-Phenyl-furan-3-carbonyl]-amino}-phenyl)-acetic acid
10 (10);
(4-{[5-(4-Difluoromethoxy-phenyl)-furan-3-carbonyl]-amino}-phenyl)-acetic acid (13);
(4-{[5-(4-Difluoromethoxy-phenyl)-2-methyl-furan-3-carbonyl]-amino}-phenyl)-acetic acid (18);
15 4-{[(5-Methyl-2-phenyl-furan-3-carbonyl)-amino]-methyl}-benzoic acid (20);
2-Methyl-5-phenyl-furan-3-carboxylic acid [4-(2-benzenesulfonylamino-2-oxo-ethyl)-phenyl]-amide (21);
5-(4-Methoxy-phenyl)-2-trifluoromethyl-furan-3-carboxylic
20 acid [4-(2-benzenesulfonylamino-2-oxo-ethyl)-phenyl]-amide (22);
5-(4-Methoxy-phenyl)-2-methyl-furan-3-carboxylic acid [4-(2-benzenesulfonylamino-2-oxo-ethyl)-phenyl]-amide (23);
5-Phenyl-furan-3-carboxylic acid {4-[2-oxo-2-(toluene-2-sulfonylamino)-ethyl]-phenyl}-amide (24);
25 5-(4-Difluoromethoxy-phenyl)-2-methyl-furan-3-carboxylic acid [4-(2-benzenesulfonylamino-2-oxo-ethyl)-phenyl]-amide (25);
5-(4-Difluoromethoxy-phenyl)-2-methyl-furan-3-carboxylic
30 acid {4-[2-oxo-2-(toluene-2-sulfonylamino)-ethyl]-phenyl}-amide (26);
5-(4-Difluoromethoxy-phenyl)-2-methyl-furan-3-carboxylic acid {4-[2-oxo-2-(propane-1-sulfonylamino)-ethyl]-phenyl}-

amide (27);

5-(4-Difluoromethoxy-phenyl)-2-methyl-furan-3-carboxylic acid {4-[2-(3,5-dimethyl-isoxazole-4-sulfonylamino)-2-oxo-ethyl]-phenyl}-amide (28);

5 5-(4-Difluoromethoxy-phenyl)-2-methyl-furan-3-carboxylic acid {4-[2-oxo-2-(thiophene-2-sulfonylamino)-ethyl]-phenyl}-amide (29);

5-(4-Difluoromethoxy-phenyl)-2-methyl-furan-3-carboxylic acid {4-[2-(5-methyl-pyridine-2-sulfonylamino)-2-oxo-ethyl]-phenyl}-amide (30);

5-(4-Difluoromethoxy-phenyl)-2-methyl-furan-3-carboxylic acid [4-(2-oxo-2-phenylmethanesulfonylamino-ethyl)-phenyl]-amide (31);

15 5-(4-Difluoromethoxy-phenyl)-2-methyl-furan-3-carboxylic acid {4-[2-oxo-2-(2-trifluoromethyl-benzenesulfonylamino)-ethyl]-phenyl}-amide (32);

5-(4-Difluoromethoxy-phenyl)-2-methyl-furan-3-carboxylic acid {4-[2-(4-hydroxy-benzenesulfonylamino)-2-oxo-ethyl]-phenyl}-amide (35); and

20 5-(4-Difluoromethoxy-phenyl)-2-methyl-furan-3-carboxylic acid {4-[2-(2-fluoro-benzenesulfonylamino)-2-oxo-ethyl]-phenyl}-amide (36).

The selectivity of the compound for antagonising EP₄ receptors over the other EP receptors (i.e. EP₁, EP₂, EP₃) can be quantified by dividing the K_i for EP₄ (see below) by the K_i for the other EP receptors (see below). The resulting ratio is preferably 10 or more, more preferably 100 or more.

Synthesis Examples

General Experimental Details

All reactions were carried out under an inert atmosphere of nitrogen.

5

Where products were purified by flash chromatography the stationary phase used was silica gel for chromatography, 0.035 to 0.070 mm (220 to 440 mesh) (e.g. Fluka silica gel 60). An applied pressure of nitrogen of ~10 psi was used to accelerate column elution. Thin layer chromatography (TLC) was carried out on aluminium foil plates coated with silica gel containing a fluorescent indicator (254 nm) (e.g. Fluka 60778).

10

15 Petroleum ether refers to that fraction with a boiling point of 40-60°C.

Organic solutions were dried over magnesium sulphate unless otherwise specified.

20

PS-TsCl refers to Polystyrene scavenger resin (loading 1.97 mmol/g) - Argonaut Technologies (P/N 800277)

25 *Preparative HPLC System*

Preparative HPLC was carried out on a C18-reverse-phase column (10 x 2.1 cm i.d Genesis column with 7 µm particle size), eluting with a gradient of acetonitrile (containing 0.1% trifluoroacetic acid) in water (containing 0.1% trifluoroacetic acid) at a flow rate of 5 ml/min. The gradient was started at 50% acetonitrile, and was increased at a rate of 1% per minute up to 90% acetonitrile/water unless otherwise stated. UV detection at 230 nm was used unless otherwise stated.

30

LC/MS Systems

The Liquid Chromatography Mass Spectroscopy (LC/MS) systems used are as follows.

5 LC/MS System A:

Mass Spectrometer - Platform LC with electrospray source operating in positive and negative ion mode. HP1100 system running at 2.0 mL/min, 200 µL/min split to the ESI source with inline HP1100 DAD detection and
10 SEDEX ELS detection.

Mobile Phase

A) Water 0.1 % Formic Acid

B) acetonitrile 0.1% Formic Acid

15

Gradient

Time (min)	Flow (mL/min)	%A	%B
0.00	2.0	95	5
0.50	2.0	95	5
4.50	2.0	5	95
5.00	2.0	5	95
5.50	2.0	95	5

Column - Luna 3u C18(2) 30x4.6mm

20 LC/MS System B:

Mass Spectrometer - Platform II with electrospray source operating in negative ion mode. HP1100 system running at 2.0 mL/min, 200 µL/min split to the ESI source with inline HP1100 DAD detection and SEDEX ELS
25 detection.

Mobile Phase

A) Water 0.1 % Diethylamine

B) acetonitrile

5

Gradient

Time (min)	Flow (mL/min)	%A	%B
0.00	2.0	95	5
0.50	2.0	95	5
4.00	2.0	5	95
4.50	2.0	5	95
5.00	2.0	95	5
20.00	2.0	95	5

Column - XTerra MS C18 3.5µm 4.6 x 30mm

LCMS System C:

10

Mass Spectrometer - Finnigan TSQ700 with electrospray source operating in negative ion mode.

HP1050 system running at 2.0 mL/min, 200 µL/min split to the ESI source with inline HP1050 Single wavelength UV detector at 254 nm.

15

Mobile Phase

A) Water 0.1 % Diethylamine

B) acetonitrile

Gradient

Time (min)	Flow (mL/min)	%A	%B
0.00	2.0	95	5
1.00	2.0	95	5
15.00	2.0	5	95
17.00	2.0	5	95

18.00	2.0	95	5
20.00	2.0	95	5

Column - XTerra MS C18 3.5 μ m 4.6 x 30mm

LC/MS System D:

5 Mass Spectrometer - Finnigan TSQ700 with electrospray
source operating in positive or negative ion mode.
HP1050 system running at 2.0 mL/min, 200 μ L/min split
to the ESI source with inline HP1050 Single Wavelength
10 UV detector at 254 nm.

Mobile Phase

A) Water 0.1 % formic Acid
B) acetonitrile 0.1% formic Acid


15 Gradient

Time (min)	Flow (mL/min)	%A	%B
0.00	2.0	95	5
1.00	2.0	95	5
15.00	2.0	5	95
17.00	2.0	5	95
18.00	2.0	95	5
20.00	2.0	95	5

Column - Higgins Clipius C18 5 μ m 100 x 3.0mm

Example 1: Synthesis of [(alkyl-phenyl-furan-3-carbonyl)-amino]-phenyl-acetic acids

(a) {4-[(2-Methyl-5-phenyl-furan-3-carbonyl)-amino]-phenyl}-acetic acid (2)



(i) Diisopropylethylamine (427mg) was added to a stirred solution of 2-methyl-5-phenyl-furan-3-carboxylic acid (334mg, 1.6 mmol) and ethyl-4-aminophenyl acetate (296 mg, 1.65mmoles) in N,N-dimethylformamide (30 ml). O-(7-Azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (628mg, 1.65mmoles) was added and the solution was stirred at room temperature for 18 hours. The solvent was evaporated, the residue was dissolved in dichloromethane and washed with water, 10% aqueous sodium carbonate, 1M aqueous hydrochloric acid and finally dried (MgSO₄). After evaporation of the solvent, the residue was triturated with cyclohexane and dried to afford {4-[(2-Methyl-5-phenyl-furan-3-carbonyl)-amino]-phenyl}-acetic acid ethyl ester (1) (466 mg) as a gum. LC/MS System D; R_t = 10.64mins, m/z (ES⁺) = 364 (M+H for C₂₂H₂₁NO₄).


(ii) A solution of sodium hydroxide (150mg) in water (5ml) was added to a stirred solution of {4-[(2-methyl-5-phenyl-furan-3-carbonyl)-amino]-phenyl}-acetic acid ethyl ester (1) (150mg, 0.41mmoles) in ethanol (20ml) and the mixture was stirred at room temperature for 1 hour. The solvent was evaporated and the residue was diluted with water (10ml) and

acidified to pH2 with 1M aqueous hydrochloric acid. The precipitate was collected, washed with water and the residue was triturated with cyclohexane. Recrystallisation from isopropanol afforded compound (2) (135mg) as a white solid.

5 LC/MS System C; $R_t = 4.06\text{mins}$, m/z (ES^-) = 334 (M-H for $\text{C}_{20}\text{H}_{17}\text{NO}_4$).

(b) (4-([5-(4-Methoxy-phenyl)-2-trifluoromethyl-furan-3-carbonyl]-amino)-phenyl)-acetic acid (4)

10




(i) In an analogous manner to example 1(a)(i), (4-([5-(4-Methoxy-phenyl)-2-trifluoromethyl-furan-3-carbonyl]-amino)-phenyl)-acetic acid ethyl ester (3) was synthesised from 5-(4-methoxy-phenyl)-2-trifluoromethyl-furan-3-carboxylic acid (160mg, 0.56mmol) and ethyl-4-aminophenyl acetate (100mg, 0.56mmoles). 190mg of the product was obtained as a gum.

20 LC/MS System A; $R_t = 4.15\text{mins}$, m/z (ES^+) = 448 (M+H for $\text{C}_{23}\text{H}_{20}\text{F}_3\text{NO}_5$).

(ii) In an analogous manner to example 1(a)(ii), compound (4), was synthesised from (4-([5-(4-methoxy-phenyl)-2-trifluoromethyl-furan-3-carbonyl]-amino)-phenyl)-acetic acid ethyl ester (3) (180mg, 0.403mmoles). The resulting precipitate was collected, washed with water and the residue was triturated with cyclohexane to afford compound

(4) (140mg) as a white solid. LC/MS System D; $R_t = 8.07$ mins, m/z (ES^+) = 420 ($M+H$ for $C_{21}H_{16}F_3NO_5$).


(c) 4-([5-(4-Methoxy-phenyl)-2-methyl-furan-3-carbonyl]-
5 amino)-phenyl)-acetic acid (6)



(i) In an analogous manner to example 1(a)(i), (4-[[5-(4-methoxy-phenyl)-2-methyl-furan-3-carbonyl]-amino]-phenyl)-acetic acid ethyl ester (5) was synthesised from 5-(4-methoxy-phenyl)-2-methyl-furan-2-carboxylic acid (250mg, 1.077mmol) and ethyl-4-aminophenyl acetate (193mg, 1.077mmoles). 80mg of the compound was obtained as a gum. LC/MS System A; R_t = 4.03mins, m/z (ES^+) = 394 ($M+H$ for $C_{23}H_{23}NO_5$).

15 (ii) In an analagous manner to example 1(a)(ii), compound (6), was synthesised from {4-[(5-phenyl-furan-3-carbonyl)-amino]-phenyl}-acetic acid ethyl ester (5) (50mg, 0.143mmoles). The resulting precipitate was collected, 20 washed with water and the residue was triturated with cyclohexane to afford compound (6) (14mg) as a white solid. LC/MS System D; R_t = 7.25mins, m/z (ES^+) = 322 ($M+H$ for $C_{19}H_{15}NO_4$).

25 (d) {4-[(5-Phenyl-furan-3-carbonyl)-amino]-phenyl}-acetic
acid (10)




(i) To a stirred solution of pyridinium tribromide (14.3g, 44.6mmoles) in acetic acid (20 ml) was added 3-furoic acid (5g, 44.6mmoles). The resulting mixture was heated at 40°C for 16 hours. The acetic acid was evaporated; the residue was dissolved in dichloromethane (50ml), washed with water (3x50ml) and dried (MgSO₄). The solvent was evaporated and the residue was purified by HPLC (gradient: 20% acetonitrile/80% water containing 0.1% trifluoroacetic acid to 80% acetonitrile/20% water at a rate of 1%/min) to afford 5-Bromo-furan-3-carboxylic acid (7) as a white solid. LC/MS System A; R_t = 2.52mins, m/z (ES⁻) = 189/191 (M-H for C₅H₃BrO₃).

(ii) In an analogous manner to example 1(a)(i), (4-[(5-bromo-furan-3-carbonyl)-amino]-phenyl)-acetic acid ethyl ester (8) was synthesised from 5-bromo-furan-3-carboxylic acid (7) (225mg, 1.18mmol) and ethyl-4-aminophenyl acetate (213mg, 1.18mmoles). 403mg of the product was obtained as a brown solid. LC/MS System A; R_t = 3.49mins, m/z (ES⁺) = 352/354 (M+H for C₁₅H₁₄BrNO₄).

(iii) A solution of {4-[(5-bromo-furan-3-carbonyl)-amino]-phenyl}-acetic acid ethyl ester (8) (200mg, 0.57mmoles), phenyl boronic acid (69.2mg, 0.57 mmoles), [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) (50mg, 0.06mmoles) and 2M aqueous cesium carbonate (1.2ml, 2.4mmoles) in toluene (20ml) was refluxed for 16 hours. The solvent was evaporated, the residue diluted with water (20ml) and then extracted with ethyl acetate (3x20ml). The combined organic layers were concentrated *in vacuo*. The residue was purified by flash chromatography (gradient elution with 90% cyclohexane / 10% ethyl acetate to 50% cyclohexane / 50% ethyl acetate) to afford {4-[(5-phenyl-furan-3-carbonyl)-amino]-phenyl}-acetic acid ethyl ester (9) (53mg). LC/MS System A; $R_t = 3.84\text{mins}$, m/z (ES^+) = $(\text{M}+\text{H})$ for $\text{C}_{21}\text{H}_{19}\text{NO}_4$.

(iv) In an analogous manner to example 1(a)(ii), compound (10), was synthesised from {4-[(5-phenyl-furan-3-carbonyl)-amino]-phenyl}-acetic acid ethyl ester (9) (50mg, 0.143mmoles). The resulting precipitate was collected, washed with water and the residue was triturated with cyclohexane to afford compound (10) (14mg) as a white solid. LC/MS System D; $R_t = 7.25\text{mins}$, m/z (ES^+) = 322 ($\text{M}+\text{H}$ for $\text{C}_{19}\text{H}_{15}\text{NO}_4$).

(e) (4-([5-(4-Difluoromethoxy-phenyl)-furan-3-carbonyl]-amino)-phenyl)-acetic acid (13)




(i) A solution of {4-[(5-Bromo-furan-3-carbonyl)-amino]-phenyl}-acetic acid ethyl ester (828mg, 2.35mmoles) (8), 4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenol (518mg, 2.35 mmoles), [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) (50mg) and 2M aqueous cesium carbonate (3.5ml, 3.5mmoles) in N,N-dimethylformamide (20ml) was heated in a microwave reactor at 100°C for 15 minutes. The solvent was evaporated, and the residue was purified by flash chromatography (gradient elution with 100% cyclohexane to 50% cyclohexane / 50% ethyl acetate) to afford (4-([5-(4-hydroxy-phenyl)-furan-3-carbonyl]-amino)-phenyl)-acetic acid ethyl ester (11) (328mg). LC/MS System A; R_t = 3.35mins, m/z (ES^+) = 366 ($M+H$ for $C_{21}H_{19}NO_5$).

(ii) To a stirred solution of (4-([5-(4-hydroxy-phenyl)-furan-3-carbonyl]-amino)-phenyl)-acetic acid ethyl ester (328mg, 0.898mmoles) (11) in N,N-dimethylformamide (15ml) was added potassium carbonate (190mg, 1.35 mmoles) and

potassium iodide (75mg, 0.449mmoles). Chlorodifluoromethane was bubbled through the solution at 80°C for 5 hours, and then discontinued. The reaction mixture was stirred at 80°C for 16 hours. The reaction was cooled to room temperature and quenched by the addition of water until no effervescence was observed. The resulting solution was extracted with ethyl acetate and the combined organic layers washed with brine, dried over MgSO₄ and concentrated *in vacuo*. The residue was purified by flash chromatography (gradient: 100% cyclohexane to 50% cyclohexane / 50% ethyl acetate) to afford (4-([5-(4-Difluoromethoxy-phenyl)-furan-3-carbonyl]-amino)-phenyl)-acetic acid ethyl ester (**12**) (60mg) as a yellow solid. LC/MS System A; R_t = 3.91mins, m/z (ES⁺) = 416 (M+H for C₂₂H₁₉F₂NO₅).

(iii) In an analagous manner to example 1(a)(ii), compound (**13**), was synthesised from (4-([5-(4-difluoromethoxy-phenyl)-furan-3-carbonyl]-amino)-phenyl)-acetic acid ethyl ester (**12**) (55mg, 0.132 mmoles). The resulting precipitate was collected, washed with water and the residue was triturated with cyclohexane to afford the compound (**13**) (41.8mg) as a white solid. LC/MS System D; R_t = 8.86mins, m/z (ES⁺) = 388 (M+H for C₂₂H₁₅F₂NO₅).

(f) (4-([5-(4-Difluoromethoxy-phenyl)-furan-3-carbonyl]-amino)-phenyl)-acetic acid (18)



5 (i) A solution of sodium hydroxide (4.5g) in water (10ml) was added to a stirred solution of 5-bromo-2-methyl-furan-3-carboxylic acid methyl ester (5g, 23mmoles) in methanol (70ml) and the mixture was stirred at room temperature for 16 hours. The solvent was evaporated and the residue was
 10 diluted with water (10ml) and acidified to pH2 with 1M aqueous hydrochloric acid. The precipitate was collected, washed with water, dried at 40°C to afford 5-Bromo-2-methyl-furan-3-carboxylic acid (14) (4g). LC/MS System A; R_t = 2.86min.

15

(ii) In an analogous manner to example 1(a)(i), {4-[(5-bromo-2-methyl-furan-3-carbonyl)-amino]-phenyl}-acetic acid ethyl ester (15) was synthesised from 5-bromo-2-methyl-furan-3-carboxylic acid (14) (1.8g, 8.78mmoles) and (4-amino-phenyl)-acetic acid ethyl ester (1.6g, 8.9mmoles). 3.08g of the product was obtained as a gum. LC/MS System A; R_t = 3.73mins, m/z (ES^+) = 367 ($M+H$ for $C_{16}H_{16}BrNO_4$).


20

(iii) A solution of (4-[(5-bromo-2-methyl-furan-3-carbonyl)-amino]-phenyl)-acetic acid ethyl ester (3g, 8.19mmoles) (15), 4-(4,4,5,5-Tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenol (1.8g, 8.19 mmoles), [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) (300mg, 10%) and 2M aqueous cesium carbonate (15ml, 15mmoles) in N,N-dimethylformamide (35ml) was heated in a microwave reactor at 100°C for 15 minutes. The solvent was evaporated, and the residue was purified by flash chromatography (gradient elution with 100% cyclohexane to 100% ethyl acetate) to afford (4-[[5-(4-hydroxy-phenyl)-2-methyl-furan-3-carbonyl]-amino]-phenyl)-acetic acid ethyl ester (16) (1.9g). LC/MS System A; $R_t = 3.54$ mins, m/z (ES^+) = 380 ($M+H$ for $C_{22}H_{21}NO_5$).

(iv) To a stirred solution of (4-[[5-(4-hydroxy-phenyl)-2-methyl-furan-3-carbonyl]-amino]-phenyl)-acetic acid ethyl ester (1.9g, 5mmoles) (16) in N, N-dimethylformamide (70ml) was added potassium carbonate (1.04g, 7.5 mmoles) and potassium iodide (416mg, 2.5mmoles). Chlorodifluoromethane was bubbled through the solution at 80°C for 5 hours, and then discontinued. The reaction mixture was stirred at 80°C for 16 hours. The reaction was cooled to room temperature and quenched by the addition of water until no effervescence was observed. The resulting solution was extracted with ethyl acetate (3x150ml) and the combined organic layers washed with brine (200ml), dried over $MgSO_4$ and concentrated *in vacuo*. The residue was purified by flash chromatography (gradient: 100% cyclohexane to 50% cyclohexane / 50% ethyl acetate) to afford (4-[[5-(4-difluoromethoxy-phenyl)-2-methyl-furan-3-carbonyl]-amino]-phenyl)-acetic acid ethyl ester (17) (694mg) as a white solid. LC/MS System A; $R_t = 4.13$ mins, m/z (ES^+) = 430 ($M+H$ for $C_{23}H_{21}F_2NO_5$).

(v) In an analogous manner to example 1(a)(ii), compound (18), was synthesised from 4-[[5-(4-difluoromethoxy-phenyl)-2-methyl-furan-3-carbonyl]-amino]-phenyl)-acetic acid ethyl ester (17) (694mg, 1.62 mmoles). The resulting precipitate was collected, washed with water and the residue was triturated with cyclohexane to afford compound (18) (643mg) as a white solid. LC/MS System D; $R_t = 9.46$ mins, m/z (ES^+) = 402 ($M+H$ for $C_{21}H_{17}F_2NO_5$).

Example 2: Synthesis of 4-[[5-Methyl-2-phenyl-furan-3-carbonyl)-amino]-methyl]-benzoic acid (20)




(i) In an analogous manner to example 1(a)(i), 4-[[5-methyl-2-phenyl-furan-3-carbonyl)-amino]-methyl]-benzoic acid ethyl ester (19) was synthesised from 5-methyl-2-phenyl-furan-3-carboxylic acid (79mg, 0.39mmol) and 4-aminomethyl benzoic acid ethyl ester (70mg, 0.39mmoles). 99mg of the product was obtained as a gum. LC/MS System A; $R_t = 3.88$ mins, m/z (ES^+) = 364 ($M+H$ for $C_{22}H_{21}NO_4$).

(ii) In an analogous manner to example 1(a)(ii), compound (20), was synthesised from 4-[[5-methyl-2-phenyl-furan-3-carbonyl)-amino]-methyl]-benzoic acid ethyl ester (19) (90mg, 0.247mmoles). The resulting precipitate was collected, washed with water and the residue was triturated with cyclohexane to afford compound (20) (51mg) as a white solid. LC/MS System D; $R_t = 6.82$ mins, m/z (ES^+) = 336 ($M+H$ for $C_{20}H_{17}NO_4$).


Example 3: Synthesis of alkyl-phenyl-furan-3-carboxylic acid [4-(2-sulphonylamino-2-oxo-ethyl)-phenyl] amides

- 5 (a) 2-Methyl-5-phenyl-furan-3-carboxylic acid [4-(2-benzenesulfonylamino-2-oxo-ethyl)-phenyl]-amide (21)




A stirred solution of {4-[(2-methyl-5-phenyl-furan-3-carbonyl)-amino]-phenyl}-acetic acid (2) (10mg, 0.030mmoles), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (5.7mg, 0.030mmoles), and 4-(N, N-dimethylamino)-pyridine (3.5mg) in dichloromethane (10ml), was treated with benzenesulphonamide (9mg, 0.060mmoles). The mixture was stirred at room temperature for 18 hours. The reaction mixture was concentrated *in vacuo* and the residue was dissolved in ethyl acetate, washed with 0.1M aqueous hydrochloric acid, and brine and finally dried (MgSO₄). After evaporation of the solvent, the residue was purified by HPLC (gradient: 20% acetonitrile/80% water containing 0.1% trifluoroacetic acid to 80% acetonitrile/20% water at a rate of 1%/min) to afford compound (21) (10mg) as a white solid. LC/MS System D; R_t = 9.98mins, m/z (ES⁺) = 475 (M+H for C₂₆H₂₂N₂O₅S).

- 25 (b) 5-(4-Methoxy-phenyl)-2-trifluoromethyl-furan-3-carboxylic acid [4-(2-benzenesulfonylamino-2-oxo-ethyl)-phenyl]-amide (22)



In an analogous way to example 3(a), compound (22) was synthesised from (4-([5-(4-methoxy-phenyl)-2-trifluoromethyl-furan-3-carbonyl]-amino)-phenyl)-acetic acid (4) (40mg, 0.095mmoles). The reaction mixture was concentrated *in vacuo* and the residue was dissolved in dichloromethane, washed with 0.1M aqueous hydrochloric acid, and brine and finally dried (MgSO₄). After evaporation of the solvent, the residue was triturated with cyclohexane and a white solid filtered off, which was purified by HPLC (gradient: 20% acetonitrile/80% water containing 0.1% trifluoroacetic acid to 80% acetonitrile/20% water at a rate of 1%/min) to afford compound (22) (12mg) as a white solid. LC/MS System D; R_t = 9.08mins, m/z (ES⁺) = 559 (M+H for C₂₇H₂₁F₃N₂O₆S).

(c) 5-(4-Methoxy-phenyl)-2-methyl-furan-3-carboxylic acid [4-(2-benzenesulfonylamino-2-oxo-ethyl)-phenyl]-amide (23)




In an analogous way to example 3(a), compound (23) was synthesised from (4-[[5-(4-methoxy-phenyl)-2-methyl-furan-3-carbonyl]-amino]-phenyl)-acetic acid (6) (100mg,

5 0.273mmoles). The reaction mixture was concentrated *in vacuo* and the residue was dissolved in dichloromethane, washed with 0.1M aqueous hydrochloric acid, and brine and finally dried (MgSO₄). After evaporation of the solvent, the residue was triturated with cyclohexane and a white solid
10 filtered off, which was then purified by HPLC (gradient: 20% acetonitrile/80% water containing 0.1% trifluoroacetic acid to 80% acetonitrile/20% water at a rate of 1%/min) to afford compound (23) (50mg) as a white solid. LC/MS System D; R_t = 8.58mins, m/z (ES⁺) = 405 (M+H for C₂₇H₂₄N₂O₆S).

15

(d) 5-Phenyl-furan-3-carboxylic acid (4-[2-oxo-2-(toluene-2-sulfonylamino)-ethyl]-phenyl)-amide (24)




In an analogous way to example 3(a), compound (24) was

20 synthesised from (4-[[5-phenyl-furan-3-carbonyl]-amino]-phenyl)-acetic acid (10) (40mg, 0.125mmoles) and replacing benzene-sulfonamide with toluene-2-sulfonamide. The reaction mixture was concentrated *in vacuo* and the residue purified by HPLC (gradient: 20% acetonitrile/80% water
25 containing 0.1% trifluoroacetic acid to 80% acetonitrile/20% water at a rate of 1%/min) to afford compound (24) (4mg) as an off-white solid. LC/MS System D; R_t = 9.85mins, m/z (ES⁺)








= 475 (M+H for C₂₆H₂₂N₂O₅S).

(e) Compounds derived from (4-([5-(4-Difluoromethoxy-phenyl)-2-methyl-furan-3-carbonyl]-amino)-phenyl)-acetic acid (18)




A stirred solution of (4-([5-(4-difluoromethoxyphenyl)-2-methyl-furan-3-carbonyl]-amino)-phenyl)-acetic acid (18) (20mg, 0.050mmoles), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (19.1mg, 0.010mmoles), and 4-(N,N-dimethylamino)-pyridine (6mg, 0.050mmoles) in dichloromethane (10ml), was treated with a sulphonamide (R^sS(=O)₂NH₂) (0.055mmoles). The mixture was stirred at room temperature for 1.5 hours (compounds 25 and 26), 2 hours (compound 31) or 16 hours (compounds 27 to 30, 32 and 36). The reaction mixture was concentrated *in vacuo* and the residue purified by HPLC (gradient: 20% acetonitrile/80% water containing 0.1% trifluoroacetic acid to 80% acetonitrile/20% water at a rate of 1%/min) to afford the desired compound.

Compound	R ^s	Yield (mg)	LC/MS R _t (mins) System D	m/z (ES ⁻)
25		20	10.51	539
26		12	10.79	553

27		12.5	9.98	505
28		37.8	10.38	558
29		33	10.22	545
30		45	9.87	554
31		38	10.57	553
32		42.8	10.78	607
36		60	10.32	557

(f) 5-(4-Difluoromethoxy-phenyl)-2-methyl-furan-3-carboxylic acid (4-[2-(4-hydroxy-benzenesulfonylamino)-2-oxo-ethyl]-phenyl)-amide (35)



- 5 (i) A solution of sodium hydroxide (116mg, 2.9mmoles) in water (10ml) was added to a stirred solution of 4-hydroxy-benzenesulphonamide (500mg, 2.9mmoles) in water (30 ml). This mixture was treated with acetic anhydride (295mg, 2.9mmoles) and stirred at room temperature for 4.5 hours.
- 10 The reaction mixture was filtered off and the resulting solid washed with water to afford acetic acid 4-sulphamoyl-phenyl ester (33) (348mg) as a yellow solid. LC/MS System A; R_t 2.06mins.
- 15 (ii) In an analogous way to example 3(e), (4-[[5-(4-difluoromethoxy-phenyl)-2-methyl-furan-3-carbonyl]-amino]-phenyl)-acetic acid (18) (50mg, 0.125mmoles) was treated with acetic acid 4-sulphamoyl-phenyl ester (33) (29.2mg, 0.137mmoles). The mixture was stirred at room temperature
- 20 for 16 hours. The reaction mixture was concentrated and the product (34) used as described in step (iii) without further

purification. LC/MS System A; R_t 3.89mins, m/z (ES^+) = 599
($M+H$ for $C_{29}H_{24}F_2N_2O_8S$).

(iii) Sodium methoxide (6.8mg, 0.125mmoles) and water (1ml)
5 were added to a stirred solution of acetic acid 4-[2-(4-[[5-(4-difluoromethoxy-phenyl)-2-methyl-furan-3-carbonyl]-amino}-phenyl)-acetylsulphamoyl]-phenyl ester (34) (74.5mg, 0.125mmoles) in methanol (10ml). The reaction mixture was stirred at room temperature for 30 minutes, concentrated and
10 the residue was purified by HPLC (gradient: 20% acetonitrile/80% water containing 0.1% trifluoroacetic acid to 80% acetonitrile/20% water at a rate of 1%/min) to afford compound (35) (18mg) as a white solid. LC/MS System D; R_t = 9.48mins, m/z (ES^-) = 555 ($M-H$ for $C_{27}H_{22}F_2N_2O_7S$).

15

Example 4: Biological Results

Binding ability to human EP receptors

Membranes were prepared from cells stably transfected with human EP receptor cDNA. In brief, cells were cultured to
20 confluency, scraped from culture flasks, and centrifuged (800 g, 8 minutes, 4°C). Cells were twice washed in ice cold homogenisation buffer containing 10 mMTris-HCl, 1 mM EDTA.2Na, 250 mM sucrose, 1 mM PMSF, 0.3 mM indomethacin, pH 7.4, homogenised and re-centrifuged as before. The
25 supernatant was stored on ice and pellets re-homogenised and re-spun. Supernatants were pooled and centrifuged at 40000g, 10 minutes, 4°C. Resultant membrane pellets were stored at -80°C until use.

30 For assay, membranes expressing human EP_4 , EP_3 , EP_2 or EP_1 receptors were incubated in Millipore (MHVBN45) plates containing assay buffer, radiolabelled [3H]PGE₂ and 0.1 to 10 000 nM concentrations of compounds. Incubations were

performed at suitable temperatures and for suitable times to allow equilibrium to be reached. Non-specific binding was determined in the presence of 10uM PGE₂. Bound and free radiolabel was separated by vacuum manifold filtration using appropriate wash buffers, and bound radiolabel was determined by scintillation counting. Constituents of each of the buffers are included in table 1 below.

The affinity or pK_i of each compound for each receptor was calculated from the concentration causing 50% radioligand displacement (IC₅₀) using the Cheng-Prusoff equation:

$$K_i = \frac{IC_{50}}{1 + \left(\frac{\text{radioligand concentration}}{\text{radioligand } K_D} \right)}$$

This approach follows that set out in Kenakin, T.P., Pharmacologic analysis of drug receptor interaction. Raven Press, New York, 2nd edition.

Table 1

Receptor		EP ₁	EP ₂	EP ₃	EP ₄
Protein / well		6.5µg	8µg	5µg	5µg
Final [³ H-PGE ₂]		3.6nM	3nM	2.5nM	1nM
Buffer	Assay	10mM MES pH6.0; 10mM MgCl ₂ ; 1mM EDTA, 3uM Indomethacin	10mM MES pH6.0; 10mM MgCl ₂ ; 1mM EDTA	10mM MES pH 6.0; 10mM MgCl ₂ ; 1mM EDTA, 100uM GTP-gamma-S	10mM MES pH6.0; 10mM MgCl ₂ ; 1mM EDTA, 3uM Indomethacin
	Wash	10mM MES pH6.0; 10mM MgCl ₂	10mM MES pH6.0; 10mM MgCl ₂	10mM MES pH 6.0; 10mM MgCl ₂	10mM MES pH6.0; 1mM EDTA

The results are presented as pK_i values in table 2 below.

Table 2

Compound	EP ₄	EP ₂	EP ₃
4	>5	<5	<5
6	>6	<5	<5
10	>6	<5.5	<5
13	>6	<5	<5
18	>6	<5.5	<5
20	>5	<5	<5
21	>7	<6	<5
22	>6	<6	<5
23	>7	<6	<5
24	>6	<7	<5
25	>7	<6	<5
26	>7	<6.5	<5
27	>6	<5	<5
28	>6	<5	<5
29	>7	<5.5	<5
30	>7	<6	<5
31	>7	<6	<5
32	>7	<5.5	<5.5
35	>7	<5.5	<5
36	>7	<6	<5

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.